

Fully differential VBF Higgs production at NNLO

Giulia Zanderighi (CERN, University of Oxford & ERC)


Based on 1506.02660, in collaboration with
Matteo Cacciari, Frédéric Dreyer, Alexander Karlberg, Gavin Salam

UCLA Radcor-Loopfest 2015

VBF Higgs production

Five good reasons to study VBF Higgs production:


1. VBF is the **largest cross-section that involves tree-level production**, and the second of all production processes (after gluon-gluon-fusion)



VBF Higgs production

Five good reasons to study VBF Higgs production:

2. It has a **distinctive signature** that involves two forward jets (tagging jets)




VBF Higgs production

CMS JHEP05(2014)104

Five good reasons to study VBF Higgs production:


3. Tagging jets allow one to better tag events and identify Higgs decays that have very large backgrounds (notably $H \rightarrow \tau\tau$ and $H \rightarrow b\bar{b}$)



VBF Higgs production

Five good reasons to study VBF Higgs production:

4. Higgs transverse momentum is non-zero at LO. Facilitates searches of invisible decay modes




VBF Higgs production

Plehn et al '01

Five good reasons to study VBF Higgs production:


5. Angular correlation of forward jets brings in sensitivity to CP properties of the Higgs and to non-SM Higgs interactions (small CP odd component is still allowed)



VBF Higgs production

Fully inclusive VBF Higgs production was known at **NNLO** in the structure function approach

Bolzoni et al '10 - '11



$\sqrt{S} = 7 \text{ TeV}$			
Higgs mass	LO	NLO	NNLO
120	$1.235^{+0.131}_{-0.116}$	$1.320^{+0.054}_{-0.022}$	$1.324^{+0.025}_{-0.024}$
160	$0.857^{+0.121}_{-0.099}$	$0.915^{+0.046}_{-0.016}$	$0.918^{+0.019}_{-0.015}$
200	$0.614^{+0.106}_{-0.082}$	$0.655^{+0.038}_{-0.012}$	$0.658^{+0.015}_{-0.010}$
300	$0.295^{+0.070}_{-0.049}$	$0.314^{+0.022}_{-0.010}$	$0.316^{+0.008}_{-0.004}$
400	$0.156^{+0.045}_{-0.030}$	$0.166^{+0.013}_{-0.007}$	$0.167^{+0.005}_{-0.001}$


The calculation suggests **tiny** renormalization/factorization scale **uncertainties** ($\sim 1\text{-}2\%$). NNLO well within the NLO band

VBF Higgs production

However, **no realistic VBF cuts** can be applied to it, as the calculation is **totally inclusive over hadronic final states that give the same vector-boson momenta**


Differential VBF Higgs production known up to now only to NLO (+PS) and also suggests small uncertainties

Figy, Oleari, Zeppenfeld '03



The structure function approach

Schematically, think of VBF as $\text{DIS} \times \text{DIS}$ with no cross-talk between radiation from the **upper** and **lower** sector (factorized approximation). Since the DIS coefficients used are inclusive over the hadronic final state, **the calculation cannot provide differential results**




Simple kinematics

Key observation:

If the scattering is Born like, then the vector boson-momenta q_i , and on-shell conditions, fix the incoming and outgoing parton momenta:

$$p_{\text{in},i} = x_i P_i \quad p_{\text{out},i} = x_i P_i - q_i \quad x_i = \frac{q_i^2}{2q_i P_i}$$




Going fully differential

This work: going beyond structure function approach.

Based on two ingredients

1. the inclusive contribution

- use the SF approach and use four-vectors q_1, q_2 to assign Born-like (i.e. $2 \rightarrow H + 2$) kinematics using the previous eqs.
- use the projected Born-like momenta to compute differential distributions



Going fully differential


This work: going beyond structure function approach.
Based on two ingredients

2. the exclusive contribution

- use the VBF H + 3 jet NLO calculation in the factorized approximation

Figy et al '07 [NLO]; Jaeger et al '14 [NLO+PS]

- keep track, for each parton, whether it belongs to upper/lower sector; this makes it possible to deduce vector-boson momenta, q_1, q_2
- for each event (weight w), add a counter-event with projected Born kinematics (weight $-w$) deduced from q_1, q_2



Going fully differential

Schematically:

$$\begin{aligned}\sigma &= \int d\Phi_B (B + V) + \int d\Phi_R R \\ &= \underbrace{\int d\Phi_B (B + V) + \int d\Phi_R R_{P2B}}_{\text{From inclusive contribution}} + \underbrace{\int d\Phi_R R - \int d\Phi_R R_{P2B}}_{\text{Finite, from exclusive contribution}}\end{aligned}$$

(P2B = Projection to Born)

Combining the two pieces:

- from the exclusive contributions we get the full contributions from double-real and one-loop single-real
- after integration over phase-space, counter-events cancel projected tree-level double real and one-loop single real contributions from the inclusive

The sum gives thus the complete, fully differential NNLO result

Going fully differential

Schematically:



$$\sigma = \int d\Phi_B (B + V) + \int d\Phi_R R$$

P2B = Projection to Born

$$= \underbrace{\int d\Phi_B (B + V) + \int d\Phi_R R_{P2B}}_{\text{From inclusive contribution}} + \underbrace{\int d\Phi_R R - \int d\Phi_R R_{P2B}}_{\text{Finite, from exclusive contribution}}$$

From inclusive contribution

Finite, from exclusive contribution



The sum gives thus the complete, fully differential NNLO result

Practicalities

For the **inclusive part** we have

- taken the phase-space from POWHEG's VBF_H
- matrix elements coded with structure functions evaluated using parametrized versions of the DIS coefficient functions
- the structure functions evaluated with the package HOPPET
<https://hoppet.hepforge.org>

Checks

- against private version of structure-function calculation ([thanks to Marco Zaro](#))
- of structure functions with APFEL 2.4.1
- approx vs exact coefficient functions (negligible difference)

Practicalities

For the **exclusive part** we have

- taken the VBF_HJJJ calculation in POWHEG
- extended POWHEG's tags to uniquely associate radiation with each sector
- for each event, uniquely determined the vector-boson momenta q_1 , q_2 and hence the counter-event (with weight $-w$)


Checks

- results for VBF_HJJJ unchanged
- sum of inclusive + exclusive at NLO, agrees with VBF_H (NLO)
- once the rapidity between the two jets increases, there is a decreasing rate of partons assigned to the “wrong” sector

Check of tagging

- partons are tagged as up or down (U/D)
- classify events into 3- or 4- jet events
- check if the U/D assignment of the partons in a given jet corresponds to the jet rapidity (positive or negative)
- the rate for “non-correspondence” must decrease when the rapidity separation between the leading jets increases


similar plots
available for
gluons in opposite
side (UD,DU)



Check of tagging

- partons are tagged as up or down (U/D)
- classify events into 3- or 4- jet events
- check if the U/D assignment of the partons in a given jet corresponds to the jet rapidity (positive or negative)
- the rate for “non-correspondence” must decrease when the rapidity separation between the leading jets increases

same plot with
O(1) bug in the
virtual contribution



Phenomenology

Take **13 TeV** LHC collisions. Jets: anti- k_t with $R=0.4$. $M_H = 125$ GeV, NNPDF3.0_nnlo_as0118 (*also at LO, NLO*), standard EW parameters.

Choose as central scale (which approximates well $\sqrt{Q_1 Q_2}$)

$$\mu_0^2(p_{t,H}) = \frac{M_H}{2} \sqrt{\left(\frac{M_H}{2}\right)^2 + p_{t,H}^2}$$

Take **VBF cuts**

- at least two jets with $p_{t,j} > 25 \text{ GeV}$
- the two hardest (tagging jets) should have

$$\Delta y_{j1j2} > 4.5 \quad m_{j1j2} > 600 \text{ GeV} \quad |y_j| < 4.5 \quad y_{j1} y_{j2} < 0$$


Phenomenology

Cross-sections: inclusive and with VBF cuts

	$\sigma^{(\text{no cuts})}$ [pb]	$\sigma^{(\text{VBF cuts})}$ [pb]
LO	$4.032^{+0.057}_{-0.069}$	$0.957^{+0.066}_{-0.059}$
NLO	$3.929^{+0.024}_{-0.023}$	$0.876^{+0.008}_{-0.018}$
NNLO	$3.888^{+0.016}_{-0.012}$	$0.826^{+0.013}_{-0.014}$


- NNLO outside the NLO band
- NNLO about 5% (1%) with (without) VBF cuts
- NNLO corrections appear to make jets softer, hence fewer events pass the VBF cuts (see next plots)

Distributions: $p_{t,j1}$ and $p_{t,j2}$



- NNLO corrections appear to make jets softer
- NNLO corrections up to $\sim 10-12\%$, typically outside the NLO band


Distributions: $p_{t,H}$ and $\Delta y_{j1,j2}$



- sometimes **parton-shower (NLOPS)** agrees well with **NNLO** ($p_{t,H}$)
- sometimes it does not ($\Delta y_{j1,j2}$)
- non-trivial kinematic dependence of K-factors (NLO/LO and NNLO/NLO)

3 versus 7 scale bands for $p_{t,H}$

3 scales: black lines; 7 scales: all lines




3 scales: $\mu_R = \mu_F = \mu_0\{1/2, 1, 2\}$

7 scales: $(\mu_R, \mu_F) = \mu_0\{(1/2, 1/2), (1/2, 1), (1, 1/2), (1, 1), (1, 2), (2, 1), (2, 2)\}$

3 versus 7 scale bands for $p_{t,H}$

3 scales: black lines; 7 scales: all lines



Conclusion: 3 and 7 scale bands very similar


Conclusions

- shown first **fully differential NNLO** results for VBF Higgs production using a **new “projection to Born”** method
- NNLO reveals that practical VBF (i.e. with cuts) has non-trivial effects beyond NLO, hence **differential NNLO is necessary for precision phenomenology** (corrections up to 10-12%)
- power of the method highlighted by the fact that **NNLO has been achieved for the first time for a $2 \rightarrow 3$ LHC process** (thanks also to the fact. approx)
- this method opens up the prospect for the only $N^3\text{LO}$ hadron-collider calculation in the foreseeable future beyond $2 \rightarrow 1$


Extra slides

Different NLOPS with POWHEG

Comparison between different showers with and without hadronization within NLOPS-POWHEG. Similar effects for other observables.




Different PDFs at various orders



- LO with LO PDFs
- NLO with NLO PDFs
- NNLO with NNLO PDFs

Different PDFs at various orders



- LO with LO PDFs
- NLO with NLO PDFs
- NNLO with NNLO PDFs